

# Stable Isotope Applications in Freshwater Zooplankton Ecology: Indicators of Trophic Dynamics and Environmental Change

<sup>1</sup>\*T. Venkateswara Rao, <sup>2</sup>B. Chakravarthi, <sup>3</sup>N. Sreenivas, <sup>4</sup>T. Sushma, <sup>5</sup>C. Smyrna, <sup>6</sup>Y. Gowthami, <sup>7</sup>B. Devi

<sup>1,2,4,5,6,7</sup>Department of Zoology & Aquaculture, Pithapur Raja's Govt. College, Kakinada, Andhra Pradesh, India

<sup>3</sup>Department of Zoology, Government Degree College, Ramachandrapuram, Andhra Pradesh, India

\*Corresponding Author: Mr. T. Venkateswara Rao [vrnkytatapudi9969@gmail.com](mailto:vrnkytatapudi9969@gmail.com)

**Abstract** - Stable isotopes of carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) are pivotal in aquatic food web studies, serving as intrinsic tracers of carbon sources and trophic positioning. In freshwater ecosystems, zooplankton represent an essential intermediate trophic group, rapidly integrating isotopic signals from lower trophic levels. This mini-review critically examines the current state of knowledge regarding stable isotope applications in freshwater zooplankton research, with a focus on quantifying anthropogenic impacts such as eutrophication, land-use conversion, and climate-mediated hydrological shifts. The review synthesizes key empirical findings, methodological protocols, and statistical approaches for interpreting isotopic variability. Recent advancements in compound-specific isotope analysis (CSIA), baseline normalization, and mixing models are discussed. Emphasis is placed on tropical and subtropical regions, including South Indian aquatic systems, where isotopic frameworks remain underutilized. This synthesis highlights the potential of stable isotope techniques to elucidate biogeochemical pathways, refine ecosystem models, and support biomonitoring under accelerating global change.

**Keywords:** Stable isotopes, aquatic ecosystems, zooplankton and compound-specific isotope analysis (CSIA).

## I. Introduction

Freshwater zooplankton are central to aquatic food webs, functioning as conduits for energy transfer between primary producers and higher trophic levels (1). Their sensitivity to environmental perturbations and rapid turnover rates make them valuable bioindicators of ecosystem integrity. Stable isotope analysis (SIA) of carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) enables quantification of assimilated energy sources and trophic positioning, offering time-integrated dietary information that overcomes limitations of gut content analysis (2). Zooplankton thus serve as ideal candidates for applying isotopic tracers to investigate ecosystem-level shifts resulting from anthropogenic and climatic drivers (3).

## II. Theoretical Foundations of Stable Isotope Ecology in Zooplankton Studies

( $\delta^{13}\text{C}$ ) reflects the isotopic composition of primary producers and can distinguish between carbon derived from autochthonous (phytoplankton) versus allochthonous (terrestrial) sources (4). ( $\delta^{15}\text{N}$ ), enriched by  $\sim 3.4\text{‰}$  per trophic level, provides estimates of relative trophic positioning. Zooplankton isotopic signatures are modulated by dietary sources, assimilation efficiencies, lipid content, and physiological processes (5). Discriminating isotopic variation requires accounting for spatial-temporal heterogeneity in baseline values, often represented by suspended particulate organic matter (SPOM) or primary consumers (6).

## III. Environmental Change Indicators through SIA

### 3.1 Eutrophication

Anthropogenic nitrogen influxes increase ( $\delta^{15}\text{N}$ ) in phytoplankton and downstream consumers (7). Elevated ( $\delta^{15}\text{N}$ ) in zooplankton correlates with agricultural runoff, aquaculture discharge, and sewage input. Multi-seasonal studies reveal peak enrichment during dry seasons due to reduced dilution effects (8).

### 3.2 Land-Use Conversion

Alterations in ( $\delta^{13}\text{C}$ ) among zooplankton assemblages signal shifts in basal production sources (9). Deforestation and urbanization increase terrestrial carbon inputs, elevating ( $\delta^{13}\text{C}$ )-depleted allochthonous signatures in zooplankton (10). Comparative gradient studies have shown clear isotopic contrasts between forested catchments and peri-urban zones (11).

### 3.3 Climatic and Hydrological Regime Shifts

Climate-induced hydrological fluctuations modulate nutrient loading, residence time, and primary productivity (12). These processes alter baseline isotope signatures,

complicating trophic interpretation. In monsoon-impacted systems, temporal isotopic shifts are pronounced and require high-frequency sampling (13).

#### IV. Taxon-Specific Isotopic Dynamics

Intertaxonomic differences in isotopic ratios arise from variable trophic niches and metabolic pathways (14). Copepods, typically omnivorous or carnivorous, exhibit higher ( $\delta^{15}\text{N}$ ) than filter-feeding cladocerans (15). Rotifers, often detritivorous or bacterivorous, reflect low-trophic dietary sources. Intraspecific variability further depends on ontogeny, life-history strategies, and reproductive modes (16). These dynamics necessitate taxon-level resolution in isotope-based studies (16).

#### V. Methodological Considerations

Analytical precision in SIA depends on rigorous sample preparation. Lipid extraction or mathematical normalization is required to correct ( $\delta^{13}\text{C}$ )- depletion (17). Preservation with formalin or ethanol can introduce isotopic artifacts; freezing remains the preferred method. Sufficient biomass is essential, often necessitating pooled samples (18). Baseline characterization using SPOM, periphyton, or primary consumers enhances trophic fractionation accuracy (19).

### VI. Case Studies in South Indian Freshwater Ecosystems

#### 6.1 Kolleru Lake

Elevated ( $\delta^{15}\text{N}$ ) values in zooplankton during post-monsoon months indicate intensified aquaculture effluent input (20). Isotope biplots ( $\delta^{13}\text{C}$ ) vs ( $\delta^{15}\text{N}$ ) show clear separation between dry and wet season clusters, reflecting shifts in nutrient sourcing (21).

#### 6.2 Godavari River Basin

Longitudinal isotope transects reveal downstream enrichment in ( $\delta^{15}\text{N}$ ), aligned with increasing agricultural density (22). SPOM ( $\delta^{13}\text{C}$ ) declines concurrently, highlighting a shift from phytoplankton- to detritus-dominated carbon pathways (23).

#### 6.3 Chilika Lagoon

Brackish transition zones exhibit spatially heterogeneous ( $\delta^{13}\text{C}$ ) and ( $\delta^{15}\text{N}$ ) values (24). Zooplankton isotopic variability aligns with salinity gradients and freshwater flushing events (25).

### VII. Emerging Techniques and Analytical Frameworks

Compound-specific isotope analysis (CSIA), particularly of amino acids, enables disentanglement of trophic enrichment from baseline shifts (26). Bayesian mixing models (e.g., MixSIAR, SIAR) offer probabilistic dietary source attribution (27). Integration with remote sensing (chlorophyll-a, turbidity) and machine learning enhances spatial-temporal resolution in ecosystem diagnostics (28).

Table 1: Isotopic Indicators of Environmental Drivers in Zooplankton Studies

Environmental Gradient	Indicator Isotope	Expected Response in Zooplankton	Supporting Metric	System Example
Nutrient loading (N)	( $\delta^{15}\text{N}$ ) increase	Elevated trophic positioning	SPOM-zooplankton ( $\delta^{15}\text{N}$ ) offset	Kolleru lake
Carbon source shift	( $\delta^{13}\text{C}$ ) shift	Autochthonous vs. allochthonous	Zooplankton SPOM ( $\delta^{13}\text{C}$ ) differential	Godavari Basin
Salinity/climate Gradient	( $\delta^{13}\text{C}$ ) and ( $\delta^{15}\text{N}$ )	Taxonomic segregation	Salinity-zonation vs. isotope map	Chilika Lagoon
Temporal Nutrient pulses	( $\delta^{15}\text{N}$ ) fluctuation	Seasonal trophic plasticity	Biplot time series	Monsoon Lakes

### VIII. Limitations and Prospects

Challenges include baseline isotopic variability, limited taxonomic resolution, and constraints in tropical systems with high temporal turnover. Future studies should prioritize long-term monitoring, inclusion of isotopically distinct end-members, and multi-tracer approaches. Combining SIA with

fatty acid profiling, DNA metabarcoding, and geochemical tracers holds promise for ecosystem-wide inference.

### IX. Conclusion

Stable isotope analysis offers an incisive tool for elucidating trophic dynamics, nutrient sourcing, and ecosystem responses to environmental change in freshwater

systems. Zooplankton, due to their ecological position and physiological sensitivity, are optimal indicators. Advances in isotopic techniques and integrative frameworks promise to expand their role in ecosystem surveillance and restoration planning.

## REFERENCES

- [1] Cole JJ, Carpenter SR, Kitchell J, Pace ML, Solomon CT, Weidel B. Strong evidence for terrestrial support of zooplankton in small lakes based on stable isotopes of carbon, nitrogen, and hydrogen. *Proc Natl Acad Sci [Internet]*. 2011 Feb 1;108(5):1975–80. Available from: <https://doi.org/10.1073/pnas.1012807108>
- [2] Feuchtmayr H, Grey J. Effect of preparation and preservation procedures on carbon and nitrogen stable isotope determinations from zooplankton. *Rapid Commun Mass Spectrom [Internet]*. 2003 Dec 15;17(23):2605–10. Available from: <https://doi.org/10.1002/rcm.1227>
- [3] Hou W, Gu B, Zhang H, Gu J, Han BP. The relationship between carbon and nitrogen stable isotopes of zooplankton and select environmental variables in low-latitude reservoirs. *Limnology [Internet]*. 2013;14(1):97–104. Available from: <https://doi.org/10.1007/s10201-012-0388-2>
- [4] Schell DM, Barnett BA, Vinette KA. Carbon and nitrogen isotope ratios in zooplankton of the Bering, Chukchi and Beaufort seas. *Mar Ecol Prog Ser [Internet]*. 1998;162:11–23. Available from: <https://www.int-res.com/abstracts/meps/v162/meps162011>
- [5] Grey J, Jones RI, Sleep D. Stable isotope analysis of the origins of zooplankton carbon in lakes of differing trophic state. *Oecologia [Internet]*. 2000;123(2):232–40. Available from: <https://doi.org/10.1007/s004420051010>
- [6] Jones RI, Grey J, Sleep D, Arvola L. Stable Isotope Analysis of Zooplankton Carbon Nutrition in Humic Lakes. *Oikos [Internet]*. 1999 Aug 6;86(1):97–104. Available from: <http://www.jstor.org/stable/3546573>
- [7] Marcus L, Virtue P, Nichols PD, Meekan MG, Pethybridge H. Effects of sample treatment on the analysis of stable isotopes of carbon and nitrogen in zooplankton, micronekton and a filter-feeding shark. *Mar Biol [Internet]*. 2017; 164(6):124. Available from: <https://doi.org/10.1007/s00227-017-3153-6>
- [8] Williams RL, Wakeham S, McKinney R, Wishner KF. Trophic ecology and vertical patterns of carbon and nitrogen stable isotopes in zooplankton from oxygen minimum zone regions. *Deep Sea Res Part I Oceanogr Res Pap [Internet]*. 2014; 90:36–47. Available from: <https://www.sciencedirect.com/science/article/pii/S096706371400065X>
- [9] Kankaala P, Taipale S, Li L, Jones RI. Diets of crustacean zooplankton, inferred from stable carbon and nitrogen isotope analyses, in lakes with varying allochthonous dissolved organic carbon content. *Aquat Ecol [Internet]*. 2010; 44(4):781–95. Available from: <https://doi.org/10.1007/s10452-010-9316-x>
- [10] Lee JY, Kim JK, Owen JS, Choi Y, Shin K, Jung S, et al. Variation in carbon and nitrogen stable isotopes in POM and zooplankton in a deep reservoir and relationship to hydrological characteristics. *J Freshw Ecol [Internet]*. 2013 Mar 1;28(1):47–62. Available from: <https://doi.org/10.1080/02705060.2012.689999>
- [11] Grey J, Jones RI, Sleep D. Seasonal changes in the importance of the source of organic matter to the diet of zooplankton in Loch Ness, as indicated by stable isotope analysis. *Limnol Oceanogr [Internet]*. 2001 May 1;46(3):505–13. Available from: <https://doi.org/10.4319/lo.2001.46.3.0505>
- [12] Driscoll ZG, Bootsma HA, Christiansen E. Zooplankton trophic structure in Lake Michigan as revealed by stable carbon and nitrogen isotopes. *J Great Lakes Res [Internet]*. 2015;41:104–14. Available from: <https://www.sciencedirect.com/science/article/pii/S0380133015000908>
- [13] SYVÄRANTA J, HÄMÄLÄINEN H, JONES RI. Within-lake variability in carbon and nitrogen stable isotope signatures. *Freshw Biol [Internet]*. 2006 Jun 1;51(6):1090–102. Available from: <https://doi.org/10.1111/j.1365-2427.2006.01557.x>
- [14] Gu B, Schell DM, Alexander V. Stable Carbon and Nitrogen Isotopic Analysis of the Plankton Food Web in a Subarctic Lake. *Can J Fish Aquat Sci [Internet]*. 1994 Jun 1;51(6):1338–44. Available from: <https://doi.org/10.1139/f94-133>
- [15] Hill JM, McQuaid CD. Stable isotope methods: The effect of gut contents on isotopic ratios of zooplankton. *Estuar Coast Shelf Sci [Internet]*. 2011; 92(3):480–5. Available from: <https://www.sciencedirect.com/science/article/pii/S0272771411000436>
- [16] VENTURA M, CATALAN J. Incorporating life histories and diet quality in stable isotope interpretations of crustacean zooplankton. *Freshw Biol [Internet]*. 2008 Jul 1;53(7):1453–69. Available from: <https://doi.org/10.1111/j.1365-2427.2008.01976.x>
- [17] Koppelman R, Böttger-Schnack R, Möbius J, Weikert H. Trophic relationships of zooplankton in the eastern Mediterranean based on stable isotope measurements. *J*

- Plankton Res [Internet]*. 2009 Jun 1;31(6):669–86. Available from: <https://doi.org/10.1093/plankt/fbp013>
- [18] Mullin MM, Rau GH, Eppley RW. Stable nitrogen isotopes in zooplankton: Some geographic and temporal variations in the North Pacific. *Limnol Oceanogr [Internet]*. 1984 Nov 1;29(6):1267–73. Available from: <https://doi.org/10.4319/lo.1984.29.6.1267>
- [19] Hannides CCS, Popp BN, Landry MR, Graham BS. Quantification of zooplankton trophic position in the North Pacific Subtropical Gyre using stable nitrogen isotopes. *Limnol Oceanogr [Internet]*. 2009 Jan 1;54(1):50–61. Available from: <https://doi.org/10.4319/lo.2009.54.1.0050>
- [20] Gu B, Schelske CL, Hoyer M V. Stable isotopes of carbon and nitrogen as indicators of diet and trophic structure of the fish community in a shallow hypereutrophic lake. *J Fish Biol [Internet]*. 1996 Dec 1;49(6):1233–43. Available from: <https://doi.org/10.1111/j.1095-8649.1996.tb01792.x>
- [21] Harris BP, Young JW, Revill AT, Taylor MD. Understanding diel-vertical feeding migrations in zooplankton using bulk carbon and nitrogen stable isotopes. *J Plankton Res [Internet]*. 2014 Jul 1;36(4):1159–63. Available from: <https://doi.org/10.1093/plankt/fbu026>
- [22] Gorbatenko KM, Lazhentsev AE, Kiyashko SI. Seasonal dynamics of the trophic status of zooplankton in the Sea of Okhotsk (based on data from stable carbon- and nitrogen-isotope analysis). *Russ J Mar Biol [Internet]*. 2014;40(7):519–31. Available from: <https://doi.org/10.1134/S1063074014070025>
- [23] PERSAUD AD, DILLON PJ, LASENBY D, YAN ND. Stable isotope variability of meso-zooplankton along a gradient of dissolved organic carbon. *Freshw Biol [Internet]*. 2009 Aug 1;54(8):1705–19. Available from: <https://doi.org/10.1111/j.1365-2427.2009.02224.x>
- [24] Goering J, Alexander V, Haubenstock N. Seasonal variability of stable carbon and nitrogen isotope ratios of organisms in a North Pacific Bay. *Estuar Coast Shelf Sci [Internet]*. 1990;30(3):239–60. Available from: <https://www.sciencedirect.com/science/article/pii/0272771490900502>
- [25] Dehairs F, Bouillon S, Mohan PC, Sreenivas N. Sources of suspended organic matter and selective feeding by zooplankton in an estuarine mangrove ecosystem as traced by stable isotopes. *Mar Ecol Prog Ser [Internet]*. 2000;208:79–92. Available from: <https://www.int-res.com/abstracts/meps/v208/meps208079>
- [26] Matthews B, Mazumder A. Compositional and interlake variability of zooplankton affect baseline stable isotope signatures. *Limnol Oceanogr [Internet]*. 2003 Sep 1;48(5):1977–87. Available from: <https://doi.org/10.4319/lo.2003.48.5.1977>
- [27] Smith SL, Henrichs SM, Rho T. Stable C and N isotopic composition of sinking particles and zooplankton over the southeastern Bering Sea shelf. *Deep Sea Res Part II Top Stud Oceanogr [Internet]*. 2002;49(26):6031–50. Available from: <https://www.sciencedirect.com/science/article/pii/S0967064502003326>
- [28] Montoya JP, Carpenter EJ, Capone DG. Nitrogen fixation and nitrogen isotope abundances in zooplankton of the oligotrophic North Atlantic. *Limnol Oceanogr [Internet]*. 2002 Nov 1;47(6):1617–28. Available from: <https://doi.org/10.4319/lo.2002.47.6.1617>

#### Citation of this Article:

T. Venkateswara Rao, B. Chakravarthi, N. Sreenivas, T. Sushma, C. Smyrna, Y. Gowthami, & B. Devi. (2025). Stable Isotope Applications in Freshwater Zooplankton Ecology: Indicators of Trophic Dynamics and Environmental Change. *International Research Journal of Innovations in Engineering and Technology - IRJIET*, 9(10), 9-12. Article DOI <https://doi.org/10.47001/IRJIET/2025.910002>

\*\*\*\*\*